
Structural Cross-section of Strike Line E




Salinity class and lithology interpretations for the Yegua, Sparta, Queen City, Carrizo, and Wilcox aquifers, Central Texas




A portion of the geophysical well log from well 15234 between 90 and 1230 feet below ground surface. The left track of the log shows the Spontaneous Potential curve with superimposed estimated salinity class The center track shows depth below ground surface in units of feet. The right track shows the shallow and deep resistivity curves (solid and dashed lines, respectively) with superimposed lithology interpretations The far right track shows the induction conductivity curve.

Geoscientist Seal The contents of this report (including figures, tables, and plates) document the work of the following licensed Texas geoscientists: John E. Meyer, P.G., No. 2026 Mr. Meyer was responsible for working on all aspects of the study and John E. Meyer preparing the report. The seal appearing on this document was authorized on Geology No.2026



Alysa Suydam, P.G., No. 15118

Ms. Suydam interpreted stratigraphy, lithology, and total dissolved solids from geophysical well logs, interpolated net sand GIS rasters, delineated salinity classes, calculated groundwater volumes, created report figures, and prepared cross-sections. Ms. Suydam completed this work as a G.I.T. under the direct supervision of Mr. Meyer and Ms. Croskrey. The seal appearing on this document was authorized on December 28, 2020.



The aquifers mapped by the Brackish Resources Aquifer Characterization System (BRACS) team at the Texas Water Development Board (TWDB) in Brackish Groundwater in Aquifers of the Upper Coastal Plains, Central Texas (Meyer and others 2020), are the Wilcox, Carrizo, Queen City, Sparta, and Yegua aquifers (listed oldest to youngest). The team mapped these aquifers in all or parts of 14 counties (Atascosa, Bastrop, Bexar, Caldwell, Dewitt, Fayette, Gonzales, Guadalupe, Karnes, Lavaca, Lee, Live Oak, Williamson, and Wilson counties), five regional water planning areas (G, K, L, P, and N), and nine groundwater conservation districts.

BRACS studies provide Texans with an estimate of the location and quantity of brackish groundwater, as groundwater salinity is an important parameter for desalination. Groundwater salinity classes are mapped as fresh (0-999 mg/L TDS), slightly saline (1,000-2,999 mg/L TDS), moderately saline (3,000-9,999 mg/L TDS), very saline (10,000-34,999 mg/L TDS), brine (greater than or equal to 35,000 mg/L TDS), or some combination of these classes (Winslow and Kister, 1956). The BRACS team accomplishes this goal by:


- mapping a stratigraphic framework from geophysical well logs,
- estimating saturated pore space using lithology interpreted from geophysical well logs and static water level,
- calculating total dissolved solids from geophysical well logs where no measured water quality samples
- delineating the extent of salinity classes based on the measured and calculated total dissolved solids, and
- calculating an estimate of in place groundwater volume per aquifer salinity class.

For Brackish Groundwater in Aquifers of the Upper Coastal Plains, Central Texas, geophysical well logs were used to make 4,652 stratigraphic picks and 5,139 groundwater salinity calculations. More than 2,000 wells with geophysical well logs or driller's descriptions assigned lithologic intervals (Figure 2). Data mining and aquifer determination yielded 3,862 measured water quality samples. All this data is interrelated and provided the foundation to map and characterize the groundwater of the study area.

GIS datasets from this study, for example formation surface elevation rasters and net sand point value shapefiles, can be downloaded from the Texas Water Development Board's website:


http://www.twdb.texas.gov/innovativewater/bracs/ studies/UCP/index.asp.

In addition to the study report and GIS datasets stratigraphic, lithologic, and salinity interpretations are saved in the BRACS Database. It may be downloaded accompanying data dictionary: http:// www.twdb.texas.gov/innovativewater/bracs/ database.asp.



Andrea Croskrey, P.G., No. 11929 Ms. Croskrey was responsible for working on all aspects of the study and preparing the report. The seal appearing on this document was authorized on December 28, 2020.

Nathanial van Oort under the direct supervision of Mr. Meyer and Ms. Croskrey, collected well information, interpreted stratigraphy and lithology from geophysical well logs, prepared stratigraphic surface GIS rasters, and prepared report figures.



We constructed nine regional cross-sections, six strike- saline. oriented and three dip-oriented (Figure 3), to illustrate the stratigraphy, lithology, and salinity interpretations for selected wells in the project. Structural cross-section of Strike Line E (Figure 1) was constructed from Brackish Groundwater in Aquifers of the Upper Coastal Plains, Central Texas data and interpretations stored in the BRACS Database. Each well on the line is labeled with the owner's name and either the BRACS Database well ID (5 digit, auto-assigned number) or the Groundwater Database State Well Number (SWN ##-##-###). Well intervals are displayed in feet relative to mean sea level with a vertical exaggeration of 80x. An approximate ground surface is shown for illustrative purposes.

This strike-oriented line was selected to highlight groundwater salinity class mapping inthe Yegua Aquifer outcrop. Seven of eleven wells on Strike Line E display an interval of interpreted moderately saline Yegua Aguifer groundwater. However, all three measured water quality samples from the Yegua Aquifer are fresh groundwater samples. The aquifer underlying the Yegua Aquifer is the Sparta Aquifer. The Sparta Aquifer is brackish or very saline on Strike Line E. Like the Sparta Aguifer, the underlying Queen City Aguifer is also brackish or very saline. Beneath the Queen City Aquifer is the Carrizo Aquifer. The Carrizo Aquifer is primarily slightly saline on Strike Line E, whereas it was mostly fresh on strike lines B, C, and D. This leaves the Carrizo Aquifer bracketed by more saline waters above and below. Like the Sparta and Queen City aquifers above it, the Wilcox Aquifer on Strike Line E is brackish and very

Additional information and cross-sections from Brackish Groundwater in Aquifers of the Upper Coastal Plains, Central Texas (Meyer and others, 2020) are available to download from the study's webpage.

Meyer, J.E., Croskrey, A.D., Suydam, A.K., and van Oort, N., 2020, Brackish Groundwater in Aquifers of the Upper Coastal Plains, Central Texas: Texas Water Development Board Report No 385, 278 p. and 9

TWDB (Texas Water Development Board), 2019a, BRACS Database: Texas Water Development Board

TWDB (Texas Water Development Board), 2019b, Groundwater Database: Texas Water Development Board.

Winslow, A.G., and Kister, L.R., 1956, Saline-water resources of Texas: U.S. Geological Survey Water-Supply Paper 1365, 105 p.




Figure 3. Location of cross-section lines relative to the study area, formation outcrops, and Texas counties. BRAC (Brackish Resources Aquifer Characterization System) well point label is the well ID in the BRACS Database GWDB (Groundwater Database) well point label is the state well number in the GWDB Database